

AD-A042 708

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER
NONNEUTRAL PLASMA EXPANSION.(U)

F/G 20/9

MAY 77 K E LONNGREN
MRC-TSR-1747

DAAG29-75-C-0024
NL

UNCLASSIFIED

1 OF 1

ADAD42-708



END
DATE
FILMED
9-77
DDC

ADA 042708

MRC Technical Summary Report #1747

NONNEUTRAL PLASMA EXPANSION

K. E. Lonngren

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

May 1977

(Received April 25, 1977)



Approved for public release
Distribution unlimited

Sponsored by

U.S. Army Research Office
P.O. Box 12211
Research Triangle Park
North Carolina 27709

and

National Science Foundation
Washington, D.C. 20550

DDC FILE COPY

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

NONNEUTRAL PLASMA EXPANSION

K. E. Lonngren *

Technical Summary Report #1747

May 1977

ABSTRACT

ACCESSION for	
NTIS	Section <input checked="" type="checkbox"/>
DDC	B.I. Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	SP-CIAL
A	

The self similar expansion of nonneutral plasmas is examined using various models. Analytical solutions are obtained for: (1) The expansion of a Maxwellian electron cloud governed by Ohm's Law; (2) The expansion of an electron cloud using a mobility model; (3) The expansion of an electron cloud using a cold plasma model; and (4) The expansion of a charge particle cloud with a temporally decaying nonlinear diffusion coefficient.

AMS(MOS) Subject Classification: 78.35

Key Words: Self similar solution of nonneutral plasma expansion

Work Unit Number 3 - Applications of Mathematics

* On leave from the University of Iowa.

Sponsored by

the United States Army under Contract No. DAAG29-75-C-0024 and
the National Science Foundation Grant #ENG 76-15645.

NONNEUTRAL PLASMA EXPANSION

K. E. Lonngren*

In a previous report,⁽¹⁾ we presented a methodical procedure for obtaining the self similar variables using some of the techniques of Lie Group Theory. General physical phenomena which occur in plasmas and could be modeled with equations amenable to a self-similar treatment were presented along with their solutions. An extensive list of references was also given. The purpose of this report is to summarize a pot-pourri of further examples which fall under the umbrage of "Nonneutral Plasma Expansion".

With the increased interest in relativistic electron beam fusion devices, considerable attention has been given to the study of nonneutral plasmas. The recent monograph by Davidson⁽²⁾ summarizes the past work and motivates our interest in examining various aspects of this problem. In this report, we shall examine three sets of fluid equation models and using the technique of "Self Similar Solution of Partial Differential Equations," obtain analytical solutions.

In addition, a recent experiment on the Wisconsin Multipole suggests that the cross-field diffusion coefficient can be modeled with a one-dimensional diffusion equation where the diffusion coefficient is nonlinear and exponentially decaying in time.⁽³⁾ A self-similar solution of this problem shall also be given.

In Section II, we describe the various models and present the self-similar solutions. The models are: (1) The expansion of a Maxwellian electron cloud governed by Ohm's Law; (2) The expansion of an electron cloud using a mobility model; (3) The expansion of an electron cloud using a cold plasma model; and (4) The expansion of a charge particle cloud with a temporally decaying nonlinear diffusion coefficient. Section III is the conclusion.

* On leave from the University of Iowa.

II. Various Problems

To analyze the problems, we follow the procedure given in reference 1 and only describe the physical phenomena, list the PDE, the self similar variables, the ODE and the solution to the ODE without repeating the details of the procedure in each case.

1) Expansion of a Maxwellian electron cloud governed by Ohm's Law ⁽⁴⁾

In this example, we examine the expansion into a vacuum of a thermalized electron cloud, described by an isothermal Maxwellian distribution. It is assumed that collisions are sufficiently frequent such that we can speak of a conductivity for the medium.

The governing equations are:

I) Equation of continuity

$$j_x + \rho_t = 0 \quad (1)$$

II) Ohm's Law

$$j = -\sigma \Phi_x \quad \text{and} \quad (2)$$

III) Maxwellian electrons

$$\rho = n_0 q \exp[q\Phi/k_B T_e] \quad (3)$$

where all symbols are standard. By differentiating (3) with respect to x and substituting (2), we obtain

$$j = -\sigma \frac{k_B T_e}{q} \frac{1}{\rho} \rho_x \quad (4)$$

In normalized units, (4) and (1) lead to

$$\left(\frac{1}{\psi} \psi_y\right)_y - \psi_\tau = 0 \quad (5)$$

where

$$\psi = \rho/n_0 q, \quad y = x/\lambda_D, \quad \tau = t/(\epsilon_0/\sigma)$$

and λ_D is the Debye length and ϵ_0/σ is a relaxation time.

We find that (5) admits a self-similar solution of the form

$$\eta(\xi) = \tau \psi(y, \tau), \quad \xi = y/\tau \quad (6)$$

The solution satisfies the conservation law that electron charge is conserved in space.

The boundary conditions which are germane to this problem are:

- a) $\psi(0, \tau) = \psi_0/\tau \Rightarrow \eta(\xi=0) = \psi_0$
 b) $\psi(\infty, \tau) = 0 = \psi(y, 0) \Rightarrow \eta(\xi=\infty) = 0$ i.e. "consolidation". (7)

Substituting (6) into (5), we obtain

$$\left[\frac{1}{\eta} \eta_{\xi}\right]_{\xi} = -[\xi \eta]_{\xi} \quad (8)$$

The first integral of (8) is

$$-\xi \eta + C_1 = \frac{1}{\eta} \eta_{\xi} \quad (9)$$

We shall further impose the condition that the current $j \rightarrow 0$ as $y \rightarrow \infty$. From (4), this transforms to $\frac{1}{\eta} \eta_{\xi} \rightarrow 0$ as $\xi \rightarrow \infty$. Using this and (7b), $C_1 = 0$.

The integral of (9) is

$$\eta = \frac{1}{\xi^2/2 + C_2} \quad (10)$$

The constant C_2 is determined from (7a) to be $C_2 = 1/\psi_0$. In terms of y and τ , the final result is

$$\psi = \frac{\rho}{n_0 q} = \frac{1}{\tau[y^2/2\tau^2 + 1/\psi_0]} \quad (11)$$

Charge is conserved as is shown below.

$$\int_0^{\infty} \frac{\rho}{n_0 q} dy = \frac{1}{n_0 q} \int_0^{\infty} \frac{d\xi}{[\xi^2/2 + 1/\psi_0]} = \frac{\pi}{n_0 q} \sqrt{\frac{\psi_0}{2}} = \text{const.} \quad (12)$$

In conclusion, we have examined the expansion of an electron cloud in a vacuum. Under conditions where Coulomb forces can be neglected, this calculation could model an electron cloud expansion in a plasma in time scales short with respect to ion motion.

2) Expansion of an electron cloud using a mobility model ⁽⁵⁾

Recently, considerable attention has been given to the problem of the transient behavior of the bulk electric field and space charge distribution in semi-conductors and in the conduction in dielectric and insulating fluids. It has been found prudent to use a

mobility model where the velocities of charge carriers injected from an emitting electrode are proportional to the electric field through their mobilities and the electric field is related to the charge densities of the carriers through Gauss's law. Many and Rakavy⁽⁶⁾ and Helfrich and Mark⁽⁷⁾ were probably the first to suggest that the problem could be modeled with the set of dimensionless equations

$$\begin{aligned} E_x &= \rho \\ i_x + \rho_t &= 0 \\ i &= \rho E \end{aligned} \tag{13}$$

which are Poisson's equation, the equation of continuity and a mobility definition for current respectively. The subscript x and t denote a partial differentiation with respect to space and time.

In their original paper, Many and Rakavy⁽⁶⁾ obtained a solution to (13) by looking for the characteristics of the problem. Subsequently, this approach was extended by Batra, Schechtman and Seki,⁽⁸⁾ Zahn, Tsang and Pao,⁽⁹⁾ de Oliveira and Ferreira⁽¹⁰⁾ and others. An extensive list of relevant experimental observations is given in reference 9.

As the problem is extremely important, we suggest an alternative technique for solution which will describe the spatial and temporal evolution of: I) a fixed electric field and II) a constant source of current which are both governed by (13). The technique that we shall apply is to find the "self-similar solution" of this set of partial differential equations.

Equation (13) can be written as

$$EE_{xx} + (E_x)^2 + E_{xt} = 0. \tag{14}$$

The self similar variables are

$$\phi = \frac{E}{t^{\alpha/\gamma}} \quad \text{and} \quad \xi = \frac{x}{t^{\beta/\gamma}} \tag{15}$$

where α/γ and β/γ are constants which will be specified by invariance and conservation requirements. Invariance specifies that $\alpha - \beta = -\gamma$.

Substituting (15) in (14), we write

$$(\phi - \frac{\beta}{\gamma} \xi) \phi_{\xi\xi} + (\frac{\alpha}{\gamma} - \frac{\beta}{\gamma}) \phi_{\xi} + (\phi_{\xi})^2 = 0 . \quad (16)$$

We shall obtain solutions for (16) subject to: I) A fixed electric field and II) a constant current at $x = 0$ requirement.

I) Electric field is constant at $x = 0$. We choose:

$$\alpha = 0$$

$$\beta = \gamma .$$

With these values, $\xi = x/t$ and $\phi = E$. The integral of (16) with these constants is

$$\phi \phi_{\xi} - \xi \phi_{\xi} = k_1 \quad (17)$$

where k_1 is a constant of integration. The constant is set equal to zero since $E(x = 0, t) = 0$ in order to satisfy space charge limited conditions. This specifies $\phi(\xi = 0) = 0$. The solution of (17) is $\phi = \xi$ from which we compute that

$$E = \frac{x+x_0}{t+t_0}$$

$$\rho = \frac{1}{t+t_0} \quad (18)$$

$$i = \frac{(x+x_0)}{(t+t_0)^2}$$

where the constants x_0 and t_0 have been explicitly included since (13) is invariant to translation.

II) Current is constant at $x = 0$. We choose:

$$\frac{2\alpha}{\gamma} = \frac{\beta}{\gamma} .$$

With this choice, we have $\xi = \frac{x}{t}$ and $\phi = \frac{E}{t}$ where ϕ satisfies (16) which becomes

$$(\phi - 2\xi) \phi_{\xi\xi} - \phi_{\xi} + (\phi_{\xi})^2 = 0 . \quad (20)$$

This can be integrated once to

$$(\phi - 2\xi)\phi_\xi + \phi = k_2 \quad (21)$$

The constant of integration k_2 is set equal to zero since we require that $E(x = 0, t)$ be zero for space charge limited conditions which specifies $\phi(\xi = 0) = 0$. The integral of (21) is

$$\phi = \xi \quad (22)$$

from which we compute that

$$\begin{aligned} E &= \frac{x+x_0}{t+t_0} \\ \rho &= \frac{1}{t+t_0} \\ i &= \frac{x+x_0}{(t+t_0)^2} \end{aligned} \quad (23)$$

where again the constants x_0 and t_0 have been introduced since (13) is invariant to translation. Note that this is identical to (18).

In (21), we can also obtain the solution for a non-space charge limited condition ($E(x = 0, t) \neq 0$) by setting the constant of integration k_2 equal to, say, $2i_0$. The integral of (21) can then be written as

$$\phi = \xi + i_0 \quad (24)$$

from which we compute that

$$\begin{aligned} E &= (t + t_0) \left(i_0 + \frac{x+x_0}{(t+t_0)^2} \right) \\ \rho &= \frac{1}{t+t_0} \\ i &= \frac{x+x_0}{(t+t_0)^2} + i_0 \end{aligned} \quad (25)$$

where again the constants x_0 and t_0 have been reintroduced. Note that (25) reduces to the space charge limited case for $i_0 = 0$.

In conclusion, we have shown that the set of equations which describe the Transient Space Charge Limited Current Problem admit self-similar solutions for two physically interesting boundary conditions. These solutions are valid in the initial stages before

the particles reach a second electrode which may be placed at $x = L$.

3. Expansion of an electron cloud using a cold plasma model

A model which can describe the behavior of an electron cloud expansion into a fixed ion background is to assume that Poisson's equation is an initial condition. The electrostatic approximation for the $\nabla \times \vec{B}$ Maxwell equation and the continuity equation assure that Poisson's equation is satisfied for all time. ⁽¹¹⁾

The basic equations are:

$$\begin{aligned}(\rho v)_x + \rho_t &= 0 \\ m v_t + m v v_x &= -eE \\ \nabla \times \vec{B} \sim 0 &= \epsilon_0 E_t - \rho v\end{aligned}\tag{26}$$

which are the equations of continuity and motion and Maxwell's equation respectively.

The self similar variables which satisfy the conservation law that $\int_0^\infty n dx = \text{constant}$ are

$$\epsilon = E; N = n t^2; u = v/t; \xi = x/t^2.\tag{27}$$

These are the same self similar variables that were obtained in an earlier study of the set (26) where the Ansatz that Maxwell's equation could replace Poisson's equation had not been made. ⁽¹²⁾ In the earlier study, it was not a pedestrian task to integrate the ODE.

Substituting (27) into (26), we now obtain the ODE:

$$\begin{aligned}-2N - 2\xi N_\xi + (NU)_\xi &= 0 \\ U - 2\xi U_\xi + UU_\xi &= -\epsilon \\ 2\xi \epsilon_\xi + NU &= 0.\end{aligned}\tag{28}$$

A solution of this set is

$$U = 2\xi, \quad N = 2 \quad \text{and} \quad \epsilon = -2\xi\tag{29}$$

from which we write the solution of (26) using (27) and (29) as

$$E = -\frac{2x}{t^2}, \quad n = \frac{2}{t^2}, \quad v = \frac{2x}{t}.\tag{30}$$

In conclusion, we find that in the final self similar solution, the density is

independent of position at the end rather than making it an a priori assumption in the calculation as did Gintsburg who treated a similar problem. ⁽¹³⁾

4. Expansion of charged particles with a temporally decaying nonlinear diffusion coefficient

In recent experiments on the Wisconsin Multipole, it was confirmed that the cross-field diffusion coefficient depended on time and amplitude as ⁽³⁾

$$D \sim \frac{\epsilon^{-\alpha t}}{\sqrt{n}} \quad (31)$$

Incorporating this in the one dimensional diffusion equation, we obtain

$$n_t = \left[\epsilon^{-\alpha t} \frac{1}{\sqrt{n}} n_x \right]_x \quad (32)$$

where all constants except α have been suitably normalized away. A change of variables

$$\tau = \frac{1}{\alpha} [1 - \epsilon^{-\alpha t}] \quad (33)$$

transforms (32) to

$$n_\tau = \left[\frac{1}{\sqrt{n}} n_x \right]_x \quad (34)$$

The self similar treatment of (34) is straightforward, at least for the case where the conservation law $\int_0^\infty n dx = \text{constant}$ is valid. The self similar variables are

$$N = n\tau^{2/3} \quad \text{and} \quad \xi = x/\tau^{2/3} \quad (35)$$

and the resulting ODE is

$$-\frac{2}{3}(\xi N)_\xi = \left[\frac{1}{\sqrt{N}} N_\xi \right]_\xi \quad (36)$$

If the burst of particles is symmetric at $x = 0$ such that $n_x|_{x=0} = 0$, and $n(x=0, \tau) = \tau^{-2/3}$ (36) can be integrated twice to yield

$$\sqrt{N} = \frac{2}{2 + \xi^{2/3}} \quad (37)$$

Using (33) and (35) in (37), we finally obtain

$$n(x,t) = \frac{4 \left[\frac{1}{\alpha} (1 - e^{-\alpha t}) \right]^2}{\left[2 \left(\frac{1}{\alpha} (1 - e^{-\alpha t}) \right)^{4/3} + \frac{x^2}{3} \right]^2} \quad (38)$$

We note that (38) gives a reasonably accurate qualitative description to the experimental results. This seems true even though the time scale in the experiment is sufficiently long such that normal modes have been exited.

III. Conclusion

The self similar behavior of four plasma phenomena have been described.

Acknowledgment

The author acknowledges Drs. J. Berryman, A. Hirose, G. A. Navratil, R. Post, H. Shen and M. Schonbeck for discussions of various aspects of this work.

REFERENCES

- (1) K. E. Lonngren, Proc. of NSF sponsored workshop on Plasma Physics, Ahmedabad, India, 1976, to be published in Pramana. Mathematics Research Center Technical Summary Report #1698.
- (2) R. C. Davidson, Theory of Nonneutral Plasmas, W. A. Benjamin, Inc. (1974).
- (3) J. R. Drake, J. R. Greenwood, G. A. Navratil and R. S. Post, Phys. Fluids 20, 148(1977).
- (4) K. E. Lonngren and A. Hirose, Phys. Letters, 59A, 285(1976).
- (5) K. E. Lonngren, J. Appl. Phys., to be published.
- (6) A. Many and G. Rakavy, Phys. Rev. 126, 1980(1962).
- (7) W. Helfrich and P. Mark, Z. Physik 166, 370(1962).
- (8) I. P. Batra, B. H. Schechtman and H. Seki, Phys. Rev. B 2, 1592(1970).
- (9) M. Zahn, C. F. Tsang and S. C. Pao, J. Appl. Phys. 45, 2432 (1974).
- (10) L. Nunes de Oliveira and G. F. Leal Ferreira, J. Electrostatics 1, 371(1975).
- (11) R. C. Davidson, Methods in Nonlinear Plasma Theory, Academic Press (1972), 33-34.
- (12) H. Shen and K. E. Lonngren, IEEE Trans. PS-4, 144(1976).
- (13) M. A. Gintsburg, Sov. Phys. Dokl., 19, 216(1974).

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1747	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 14 MRC-TSR-1747
4. TITLE (and Subtitle) NONNEUTRAL PLASMA EXPANSION,	5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) K. E. Lonngren	8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 ENG 76-15645	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Applications of Mathematics	
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.	12. REPORT DATE May 1977	13. NUMBER OF PAGES 9
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) UNCLASSIFIED	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U.S. Army Research Office P.O. Box 12211 Research Triangle Park North Carolina 27709 and National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Self similar solution of nonneutral plasma expansion		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The self similar expansion of nonneutral plasmas is examined using various models. Analytical solutions are obtained for: (1) The expansion of a Maxwellian electron cloud governed by Ohm's Law; (2) The expansion of an electron cloud using a mobility model; (3) The expansion of an electron cloud using a cold plasma model; and (4) The expansion of a charge particle cloud with a temporally decaying nonlinear diffusion coefficient.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

221 200